

## On the High Altitude Pseudoscorpions (Arachnida: Pseudoscorpionida) in the Old World

Petar BERON

Order Pseudoscorpionida in the World includes about 3000 species of ca. 435 living genera (429 in HARVEY, 1990). According to the latest revision of HARVEY (1992), the families are 24. Our analysis of all pseudoscorpions in the Old World found over 2200 m shows, that this altitude is reached by 158 species belonging to at least 66 genera and 16 families (Table 1).

Table 1  
Genera of Pseudoscorpions in the Old World (including Australia, New Zealand, New Caledonia and St. Helene) living at or above 2200 m

Families	Genera World	Genera > 2000 m	Genera Old World	Genera in the Old World >2200 m	>3000 m	> 4000 m
Chthoniidae	30	7	19	5	4	0
Lechytiidae	1	1	1	1	1	0
Tridenchthoniidae	16	4	9	4	3	0
Geogarypidae	3	3	3	2	1	0
Olpiidae	51	12	29	5	3	0
Gymnobisiidae	4	1	3	1	1	0
Ideonocidae	9	1	6	1	1	0
Hyidae	4	1	1	1	1	1
Neobisiidae	36	5	21	1	0	0
Syarinidae	14	1	1	1	0	0
Cheiridiidae	6	2	6	2	2	0
Sternophoridae	3	1	1	1	0	0
Atemnidae	20	6	17	6	2	0
Chernetidae	110	14	53	11	8	1
Cheliferidae	58	13	43	9	4	1
Withiidae	30	4	24	4	3	0

Only small number of pseudoscorpions live in the European mountains higher than 2000 m: 2 in the Pyrenees, 1 in Sierra Nevada, 6 in the Alps, 1 in the Apennines, 2 in the mountains of Balkan Peninsula, 12 in Caucasus. Only 4 species reach or live higher than 3000 m: *Neobisium jugorum* L. Koch (Alps, 3600 m), *N. nivale* Beier (Sierra Nevada, 3481 m), *N. anatolicum* Beier

(Caucasus, 3000 m). The only known dweller of our highest summits *Neobisium carcinoides* Hermann also goes as high as 3000 m in the Alps. Obviously, the members of *Neobisium* are monopolists on the highest parts of European mountains. Within the belt 2000-3000 m in Europe (incl. Caucasus) live also some species belonging to the genera *Chthonius* (*Ch. tetrachelatus* Preyssler in Caucasus up to 2500 m, in Iran up to 2900 m), *Roncus* (*R. microphthalmus* Daday up to 2200 m in Caucasus), *Chernes* (*Ch. montigenus* Simon up to 2740 m), from the genus *Neobisium* also *N. bernardi* Vachon (up to 2800 m), *N. delphinaticum* Beier (up to 2850 m), *N. noricum* Beier (up to 2500 m), *N. dolomiticum* Beier (up to 2400 m), etc.

The studies of Tullgren, Beier, Mahnert, Redikorzev and our own collections from Kenya, Tanzania and Uganda, have shown, that in the East and Central African mountains live at or above 2200 m at least 26 species of Pseudoscorpionida. At least 13 of them reach 3000 m, 4 - 3500 m and only *Titanatemnus palmquisti* is known to live above 4000 m.

At least 12 species of Pseudoscorpionida, belonging to eight families, have been recorded from Central Asia (above 2200 m), including 11 above 3000 m and four above 3500 m. The champions are *Bisetocreagris kaznakovi* (Redikorzev) - 4810 m (Neobisiidae, Tibet), *Dactylochelifera brachialis* Beier (4200 m, Karakorum), *Gobichelifera chelanops* (Redikorzev) (3650 m, Karakorum) and "*Chelifera*" *baltistanus* di Caporiacco (3950 m, Karakorum) (all three belonging to Cheliferidae).

In the Himalaya, over 2200 m have been recorded 29 species of Pseudoscorpionida, over 3000 m - 17, over 3500 m - eleven and over 4000 m - three (*Stenohya martensi* Schawaller - 4700 m, *Orochernes nepalensis* Beier and *Dactylochelifera macrotuberculatus* Krumpal, 4000 m, all three from Nepal). The species above 2200 m belong to 20 genera and 11 families.

The high altitude pseudoscorpions in South and North America belong to the families Chthoniidae (*Austrochthonius*), Pseudogarypidae (*Pseudogarypus*), Olpiidae (*Olpiolum*, *Progarypus*, *Serianus*, *Stenolpiodes*, *Stenolpium*), Cheliferidae (*Parachelifera*, *Haplochelifera*, *Dactylochelifera*, *Hysterochelifera*), Chernetidae (*Lustrochernes*, *Parachernes*) and Withiidae (*Parawithius*). The families Chthoniidae, Olpiidae, Cheliferidae, Chernetidae and Withiidae are common to the Old World. Pseudogarypidae occurs in North and South America and in Australia. None of the species and only two genera of Cheliferidae (*Hysterochelifera* and *Dactylochelifera*) are shared with the orael of the Old World.

Out of the remaining families, represented in the high altitude fauna of the Old World, Lechytiidae, Tridenchthoniidae, Geogarypidae, Ideoroncidae, Syarinidae, Cheiridiidae and Sternophoridae are represented in the Americas, but are not known there above 2200 m. Hyidae do not live in the Western Hemisphere. Garypidae reach high altitude in South America, but not in the Old World.

Table 2

## Species of Pseudoscorpions living at or above 2200 m in Europe, Central Asia, Himalaya and Tropical Africa

Families and genera	Europe	Central Asia	Himalaya	Tropical Africa
1	2	3	4	5
<b>Chthoniidae</b>	<b>3</b>	<b>1</b>	<b>2</b>	<b>5</b>
<i>Centrochthonius</i>	0	1	0	0
<i>Chthonius</i>	3	0	0	0
<i>Lagynochthonius</i>	0	0	1	0
<i>Tyrannochthonius</i>	0	0	1	5
<b>Lechytiidae</b>	<b>0</b>	<b>0</b>	<b>1</b>	<b>1</b>
<i>Lechytia</i>	0	0	1	1
<b>Tridenchthoniidae</b>	<b>0</b>	<b>0</b>	<b>1</b>	<b>5</b>
<i>Compsaditha</i>	0	0	0	2
<i>Ditha</i>	0	0	1	0
<i>Pycnodithella</i>	0	0	0	1
<i>Verrucadithella</i>	0	0	0	2
<b>Geogarypidae</b>	<b>0</b>	<b>2</b>	<b>2</b>	<b>4</b>
<i>Afrogarypus</i>	0	0	0	4
<i>Geogarypus</i>	0	2	2	0
<b>Olpiidae</b>	<b>0</b>	<b>2</b>	<b>1</b>	<b>1</b>
<i>Calocheiridius</i>	0	0	1	1
<i>Garypinus</i>	0	1	0	0
<i>Olpium</i>	0	1	0	0
<b>Ideoroncidae</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>1</b>
<i>Negroroncus</i>	0	0	0	1
<b>Hyidae</b>	<b>0</b>	<b>0</b>	<b>4</b>	<b>0</b>
<i>Stenohya</i>	0	0	4	0
(syn. <i>Laevigatocreagris</i> )	0	0	1	0
<b>Neobisiidae</b>	<b>17</b>	<b>2</b>	<b>2</b>	<b>1(2)</b>
<i>Bisetocreagris</i>	0	2	1	0
<i>Microbisium</i>	0	0	0	1
<i>Neobisium</i>	16	0	0	1(2)
<i>Nepalobisium</i>	0	0	1	0
<i>Roncus</i>	1	0	0	0
<b>Syarinidae</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>1</b>
<i>Ideoblothrus</i>	0	0	0	1
<b>Cheiridiidae</b>	<b>0</b>	<b>0</b>	<b>2</b>	<b>2</b>
<i>Apocheiridium</i>	0	0	1	1
<i>Cheiridium</i>	0	0	1	0

	1	2	3	4	5
<i>Cryptocheiridium</i>		0	0	0	1
<b>Atemnidae</b>		<b>0</b>	<b>1</b>	<b>2</b>	<b>7</b>
<i>Atemnus</i>		0	1	2	0
<i>Cyclatemnus</i>		0	0	0	3
<i>Micratemnus</i>		0	0	0	1
<i>Paratemnoides</i>		0	0	0	1
<i>Titanatemnus</i>		0	0	0	2
<b>Cheliferidae</b>		<b>0</b>	<b>6</b>	<b>3</b>	<b>4</b>
<i>Dactylochelifer</i>		0	5	1	0
<i>Gobichelifer</i>		0	1	0	0
<i>Hansenius</i>		0	0	0	1
<i>Chelifer</i>		0	0	0	1
<i>Hysterochelifer</i>		0	0	1	0
<i>Lophochernes</i>		0	0	1	0
<i>Microchelifer</i>		0	0	0	2
<b>Chernetidae</b>		<b>1</b>	<b>4</b>	<b>9</b>	<b>14</b>
<i>Alochernes</i>		1	1	0	0
<i>Caffrowithius</i>		0	0	0	5
<i>Ceriochernes</i>		0	0	3	0
<i>Dendrochernes</i>		0	1	1	0
<i>Lasiochernes</i>		0	0	0	1
<i>Lamprochernes</i>		0	0	1	0
<i>Megachernes</i>		0	1	3	0
<i>Nudochernes</i>		0	0	0	8
<i>Orochernes</i>		0	0	1	0
<i>Pselaphochernes</i>		0	1	0	0
<b>Withiidae</b>		<b>0</b>	<b>0</b>	<b>1</b>	<b>7</b>
<i>Ectromachernes</i>		0	0	0	1
<i>Stenowithius</i>		0	0	0	1
<i>Trichotowithius</i>		0	0	0	2
<i>Withius</i>		0	0	1	3

## Pseudoscorpionida in the Old World, known at or above 2200 m

Order PSEUDOSCORPIONIDA - up to ? 5000 m (Nepal)

### Chthoniidae - up to 3650 m (Nepal)

*Afrochthonius* Beier - up to 2590 m (*A. brincki* Beier, Lesotho), 2500 m (*A. ceylonicus* Beier, Sri Lanka)

*Centrochthonius* Beier - up to 3650 m (*C. kozlovi* Redikorzev, Nepal, Tibet)

*Chthonius* C. L. Koch - up to 3030 m (*Ch. hispanus* Beier, Sierra Nevada), 2900 m (*Ch. tetrachelatus* Preyssler, Iran, Elburz; 2500 m, Caucasus), 2400 m (*Ch. dacnodes* Navas, Spain)

*Lagynochthonius* Beier - up to 3100 m (*L. himalayensis* Morikawa, Nepal),




Fig. 1. Altitudinal distribution of Pseudoscorpionida living at and above 2200 m in the Old World. A - Chthoniidae (up to 3650 m); B - Lechytiidae (up to 3600 m); C - Tridenchthoniidae (up to 3200 m); D - Geogarypidae (up to 3200 m); E - Olpiidae (up to 3650 m); F - Hyidae (up to 5000 m); G - Ideoroncidae (up to 2250 m); H - Neobisidae (up to 5000 m); I - Syarinidae (up to 3000 m); J - Cheiridiidae (up to 3600 m); K - Sternophoridae (up to 2300 m); L - Atemnidae (up to 4100 m); M - Cheliferidae (up to 4200 m); N - Chernetidae (up to 4000 m); O - Witiidae (up to 3500 m)

2400 m (*L. tonkinensis* Beier, Vietnam; 2300 m, Thailand), 2400 m (*L. annamensis* Beier, Vietnam)

*Tyrannochthonius* Chamberlin - up to 3500 m (*T. rahmi* Beier, Nepal; *T. robustus* Beier, Sichuan, China), 3300 m (*T. brevipennis* Beier, Elgon), 3300 m (*T. meneghettii* di Caporiacco = *T. holmi* Beier, Elgon), 3025 m (*T. wittei* Beier, Kivu), 2800 m (*T. sokolovi* Redikorzev, Kenya; 2780 m, Kivu), 2550 m (*T. pachythorax* Redikorzev, Thailand), 2350 m (*T. elegans* Beier, Kivu)

#### **Lechytiidae** - up to 3600 m (Nepal)

*Lechytia* Balzan - up to 3600 m (*L. himalayana* Beier, Nepal), 2900 m (*L. maxima* Beier, Kenya)

#### **Tridenchthoniidae** - up to 3200 m (Kenya)

*Compsaditha* Chamberlin - up to 2300 m (*C. basilewskyi* Beier, Kenya), 2250 m (*C. congica* Beier, Kivu)

*Verrucadithella* Beier - up to 3200 m (*V. dilatimana* Redikorzev, Kenya), 2950 m (*V. jeanneli* Beier, Elgon)

*Pycnodithella* Beier - up to 3000 m (*P. abyssinica* Beier, Ethiopia)

*Ditha* Chamberlin - up to 3000 m (*D. proxima* Beier, Nepal), 2500 m (*D. tonkinensis* Beier, Vietnam)

#### **Geogarypidae** - up to 3200 m (Nepal, Kivu)

*Afrogarypus* Beier - up to 3200 m (*A. intermedius* Beier, Kivu), 2900 m (*A. monticola* Beier, Kenya), 2900 m (*A. zonatus* Beier, Kivu), 2200 m (*A. basilewskyi* Beier, Kenya)



*Geogarypus* Chamberlin - up to 3200 m (*G. nepalensis* Beier, Nepal), 3000 m (*G. continentalis* Redikorzev, Kirgizstan, Kungey Alatau), 2440 m (*G. angulatus* Chamberlin, India, Nilgiris), 2300 m (*G. irrugatus* Simon, Bhutan; 2200 m, Tien Shan)

**Olpiidae** - up to 3650 m (Elgon), 4100 m (Peru)

*Amblyolpium* Simon - up to 3300 m (*A. simoni* Heurtault, Tibesti, Tchad)

*Calocheiridius* Beier et Turk - up to 3650 m (*C. crassifemoratus* Beier, Elgon), 2730 m (*C. sulcatus* Beier, Nepal)

*Garypinus* Daday - up to 3000 m (*G. afghanicus minor* Beier, Afghanistan)

*Horus* Chamberlin - up to 2350 m (*H. montanus* Beier, Lesotho), 2318 m (*H. obscurus* Tullgren, Drakensberg)

*Olpium* L. Koch - up to 3000 m (*O. intermedium* Beier, Afghanistan), 2800 m (*O. tenue* Chamberlin, Tibesti, Tchad)

**Hyidae** - up to ? 5000 m (Nepal)

*Stenohya* Beier (syn. *Levigatocreagris* Curčić) - up to 5000 m (*Stenohya* sp., sub "*Levigatocreagris* / *Bisetocreagris* sp.", Nepal), 4700 m (*S. martensi* Schawaller, Nepal), 3500 m (*S. gruberi* Curčić, Nepal), 2700 m (*S. kashmirensis* Schawaller, Kashmir), 2500 m (*S. hamatus* Leclerc et Mahnert, Thailand)

**Gymnobisiidae** - up to 3203 m (South Africa)

*Gymnobisium* Beier - up to 3203 m (*G. quadrispinosum* Tullgren, South Africa)

**Ideoroncidae** - up to 2250 m (Kenya)

*Negroroncus* Beier - up to 2250 m (*N. silvicola* Mahnert, Kenya)




Fig. 2. Genera of Pseudoscorpionida living at or above 2200 m in the Old World. A - Chthoniidae (5); B - Lechytiidae (1); C - Tridenchthoniidae (4); D - Geogarypidae (2); E - Olpiidae (5); F - Hyidae (1); G - Ideoroncidae (1); H - Neobisiidae (6); I - Syarinidae (1); J - Cheiridiidae (3); K - Sternophoridae (1); L - Atemnidae (7); M - Cheliferidae (9); N - Chernetidae (11); O - Withiidae (4)

**Neobisiidae** - up to ? 5000 m (Nepal)

*Bisetocreagris* Curčić - up to ? 5000 m (*Bisetocreagris* sp., sub "*Levigatocreagris* / *Bisetocreagris* sp.", Nepal), 4810 m (*B. kaznakovi* Redikorzev, Tibet; 4000 m, Kirgizstan; 3600 m, Nepal), 2700 m (*B. tenuis* Redikorzev, Tien Shan), 2550 m (*B. indochinensis* Redikorzev, Thailand), 2400 m (*B. philippinensis* Beier, Luzon, Philippin Is.)

*Microbisium* Chamberlin - up to 3300 m (*M. dogieli* Redikorzev = *M. perpusillum* Beier, Kenya; 2300 m, Rwanda)

*Microcreagris* Balzan - up to 2200 m (*M. brevidigitata* Chamberlin, Japan)

*Neobisium* Chamberlin - up to 4100 m (*N. alticola* Beier, Anatolia), 3600 m (*N. jugorum* L. Koch, Alps), 3481 m (*N. nivale* Beier, Sierra Nevada), 3200 m (*N. kobachidzei* Beier, *N. erythroactylum* L. Koch, Caucasus), 3000 m (*N. anatolicum* Beier, Caucasus), 3203 m (*N. carcinoides* Hermann = *N. muscorum* Leach, Aberdare, Kenya; 3000 m, Alps; 2914 m, Pirin), 2850 m (*N. delfinaticum* Beier, Alps), 2800 m (*N. bernardi* Vachon, Pyrenees), 2500 m (*N. crassifemuratum* Beier, Caucasus; *N. carpaticum* Beier, Carpathes; *N. noricum* Beier, Alps; *N. fuscimanum* C. L. Koch, Caucasus; *N. validum* C. L. Koch, Caucasus; *N. labinskyi* Beier, Caucasus), 2400 m (*N. dolomiticum* Beier, Dolomiti), 2200 m (*N. ischyrum* Beier, Dolomiti)

*Nepalobisium* Beier - up to 3150 m (*N. franzi* Beier, Nepal)

*Roncus* Beier - up to 2200 m (*R. microphthalmus* Daday, Caucasus)

**Syarinidae** - up to 3000 m (Japan)

"*Orideobisium*" Kishida (nomen nudum) - up to 3000 m ("*O. takanoanum*" Kishida, nomen nudum, Japan)

*Ideoblothrus* Balzan - up to 2200 m (*I. leleupi* Beier, Kivu, sub "*Ideobisium*")




Fig. 3. Species of Pseudoscorpionida living at or above 2200 m in the Old World. A - Chthoniidae (15); B - Lechytiidae (2); C - Tridenchthoniidae (6); D - Geogarypidae (6); E - Olpiidae (7); F - Hyidae (4); G - Ideoroncidae (1); H - Neobisiidae (25); I - Syarinidae (1); J - Cheiridiidae (4); K - Sternophoridae (1); L - Atemnidae (11); M - Cheliferidae (17); N - Chernetidae (31); O - Withiidae (8)

**Cheiridiidae** - up to 3600 m (Nepal)

*Apocheiridium* Chamberlin - up to 3600 m (*A. rossicum* Redikorzev = *A. nepalense* Curčić, Nepal), 2300 m (*A. pallidum* Mahnert, Kenya)

*Cheiridium* Menge - up to 2800 m (*Ch. nepalense* Beier, Nepal)

*Cryptocheiridium* Chamberlin - up to 3200 m (*C. elgonense* Beier, Elgon), 2250 m (*C. kivuense* Beier, Kivu)

**Sternophoridae** - up to 2300 m (New Guinea)

*Afrosterophorus* Beier - up to 2300 m (*A. cavernae* Beier, Papua New Guinea)

**Atemnidae** - up to 4100 m (Kilimanjaro)

*Anatemnus* Beier - up to 2400 m (*A. angustus* Redikorzev, Vietnam)

*Atemnus* Canestrini - up to 3000 m (*A. politus* Simon, Kirgizstan; 2500 m, Karakorum; 2400 m, Kashmir; 2200 m, Nepal), 2300 m (*A. turkestanicus* Redikorzev, Bhutan)

*Cyclatemnus* Beier - up to 3000 m (*C. minor* Beier, Ethiopia), 2350 m (*C. centralis* Beier, Rwanda; *C. fallax* Beier, Elgon), 2220 m (*C. robustus* Beier, Kivu)

*Micratemnus* Beier - up to 2200 m (*M. sulcatus* Beier, Kenya)

*Oratemnus* Beier - up to 2200 m (*O. indicus* With, India)

*Paratemnoides* Harvey (pro *Paratemnus* Beier) - up to 3050 m (*Paratemnoides* sp., Kenya)

*Stenatemnus* Beier - up to 2300 m (*S. brincki* Beier, India)

*Titanatemnus* Beier - up to 4100 m (*T. palmquisti* Tullgren = *T. montanus* Beier, Tanzania, Kenya), 2900 m (*T. chappuisi* Beier, Elgon), 2780 m (*T. sjostedti* Tullgren, Kivu)




Fig. 4. Genera of Pseudoscorpionida living at or above 2500 m in the Old World. A - Chthoniidae (5); B - Lechytiidae (1); C - Tridenchthoniidae (4); D - Geogarypidae (2); E - Olpidae (4); F - Hyidae (1); G - Neobisiidae (4); H - Syarinidae (1); I - Cheiridiidae (3); J - Atemnidae (4); K - Cheliferidae (6); L - Chernetidae (9); M - Withiidae (3)



**Cheliferidae** - up to 4200 m (Karakorum)

*Amaurochelifer* Beier - up to 2400 m (*A. annamensis* Beier, Vietnam)

*Chelifer* Geoffroy - up to 2750 m (*Ch. cancroides* L., Kivu)

"*Chelifer*" - up to 3950 m ("*Ch. baltistanus*" di Caporiacco, Karakorum, nomen dubium)

*Dactylochelifer* Beier - up to 4200 m (*D. brachialis* Beier, Karakorum), 4000 m (*D. macrotuberculatus* Krumpal, Nepal), 3650 m (*D. popovi* Redikorzev, Tajikistan; 3200 m, Terskey Alatau, Kirgizstan), 2640 m (*D. monticola* Beier, Afghanistan), 2500 m (*D. redikorzevi* Beier, Kazakhstan), 2200 m (*D. vtorovi* Mahnert, Tien Shan; *D. syriacus* Beier, Syria)

*Gobichelifer* Krumpal - up to 3000 m (*G. semenovi* Redikorzev, Kirgizstan)

*Hansenius* Chamberlin - up to 2250 m (*H. kilimanjaricus* Beier, Kenya, Kilimanjaro)

*Hysterochelifer* Chamberlin - up to 3500 m (*H. nepalensis* Beier, Nepal)

*Lophochernes* Simon - up to 2600 m (*L. indicus* Beier, Nepal)

*Microchelifer* Beier - up to 2700 m (*M. granulatus* Beier, Kenya), 2300 m (*M. dentatus* Mahnert, Kenya)

*Pseudorhacochelifer* Beier - up to 2300 m (*P. schurmanni* Beier, La Palma, Canary Is.)

*Rhacochelifer* Beier - up to 3500 (*Rh. cf. subsimilis* Vachon, Tibesti), 2910 m (*Rh. anatolicus* Beier, Turkey)

**Chernetidae** - up to 4000 m (Nepal)

*Allochernes* Beier - up to 3200 m (*A. asiaticus* Redikorzev, Terskey Alatau, Kirgizstan; *Allochernes* sp., Kashmir), 2800 m (*A. tropicus* Beier, Sichuan, China), 2600 m (*A. wideri* C. L. Koch, Alps), 2200 m (*A. longipilosus* Mahnert, Tenerife)

*Caffrowithius* Beier (syn. *Plesiochernes* Vachon) - up to 3300 m (*C. elgonensis* Vachon, Elgon), 3000 m (*C. aethiopicus* Beier, Ethiopia), 2780 m (*C.*




Fig. 5. Genera of Pseudoscorpionida living at or above 3000 m in the Old World. A - Chthoniidae (4); B - Lechytiidae (1); C - Tridenchthoniidae (3); D - Geogarypidae (1); E - Olpiidae (4); F - Hyidae (1); G - Neobisiidae (4); H - Syarinidae (1); I - Cheiridiidae (2); J - Atemnidae (4); K - Cheliferidae (4); L - Chernetidae (8); M - Witihiidae (3)

*simplex* Beier, Kivu), 2300 m (*C. rusticus* Beier, Elgon; *C. calvus* Beier, Aberdare)

*Ceriochernes* Beier - up to 3200 m (*C. vestitus* Beier, Nepal), 3100 m (*C. nepalensis* Beier, Nepal), 2500 m (*C. martensi* Beier, Nepal)

*Dendrochernes* Beier - up to 3200 m (*D. cyrneus* L. Koch, Nepal; 2400 m, Kazakhstan)

*Lamprochernes* Tömösvary - up to 2350 m (*Lamprochernes* sp. ? *savignyi* Simon, Nepal)

*Lasiochernes* Beier - up to 2200 m (*L. punctiger* Beier, Kivu)

*Megachernes* Beier - up to 3650 m (*M. himalayensis* Ellingsen, Nepal; 2400 m, Kashmir), 3550 m (*M. soricicola* Beier, Nepal), 2800 m (*M. loebli* Schawaller, Nepal), 2530 m (*M. trautneri* Schawaller, Thailand), 2390 m (*M. afghanicus* Beier, Afghanistan), 2300 m (*M. papuanus* Beier, New Guinea), 2200 m (*M. limatus* Hoff et Parrack, New Guinea)

*Nudochernes* Beier - up to 3700 m (*N. crassus* Beier, Elgon; 3000 m, Aberdare), 3500 m (*N. montanus* Beier, *N. robustus* Beier, Elgon), 3130 m (*N. longipes* Beier, Mount Kenya), 3025 m (*N. gracilipes* Beier, Kivu), 3000 m (*N. nidicola* Beier, Marakwet, 2470 m, Elgon), 2900 m (*N. leleupi* Beier, Kivu), 2780 m (*N. intermedius* Beier, Kivu), 2750 m (*N. gracilimanus* Mahnert, Mount Kenya), 2600 m (*N. granulatus* Beier, Kilimanjaro), 2200 m (*N. punctiger* Beier, Kivu)

*Orochernes* Beier - up to 4000 m (*O. nepalensis* Beier, Nepal)

*Pselaphochernes* Beier - up to 3300 m (*Pselaphochernes* sp.)

*Sundochernes* Beier - up to 2550 m (*S. novaeguineae* Beier, New Guinea)

*Verrucachernes* Chamberlin - up to 2200 m (*V. montigenus* Beier, New Guinea)

## Withiidae - up to 3500 m (Elgon)

*Ectromachernes* Beier - up to 3000 m (*E. mirabilis* Beier, Ethiopia)




Fig. 6. Genera of Pseudoscorpionida living at or above 3500 m in the Old World. A - Chthoniidae (2); B - Lechytiidae (1); C - Olpiidae (1); D - Hyidae (1); E - Neobisiidae (2); F - Cheiridiidae (1); G - Atemnidae (1); H - Cheliferidae (3); I - Chernetidae (3); J - Withiidae (1)

*Stenowithius* Beier - up to 2180 m (*S. bayoni* Ellingsen = *S. ugandanus* Beier, Elgon)

*Trichotowithius* Beier - up to 3000 m (*T. abyssinicus* Beier, Ethiopia), 2400 m (*T. elgonensis* Beier, Elgon)

*Withius* Kew - up to 3500 m (*W. somalicus* Beier = *Allowithius crassus* Beier, Elgon), 3000 m (*W. abyssinicus* Beier, Ethiopia), 2500 m (*W. nepalensis* Beier, Nepal), 2300 m (*W. lewisi* Beier, Kenya)

### Pseudoscorpions living in the Old World at or above 3500 m

*Stenohya* (= *Levigatocreagris*) / *Bisetocreagris* sp. (Hyidae or Neobisiidae) - ca. 5000 m (Nepal)

*Bisetocreagris kaznakovi* (Redikorzev) (Neobisiidae) - 4810 m (Tibet)

*Stenohya* (= *Levigatocreagris*) *martensi* (Schawaller) (Hyidae) - 4700 m (Nepal)

*Dactylochelifer brachialis* Beier (Cheliferidae) - 4200 m (Karakorum)

*Titanatemnus palmquisti* (Tullgren) (= *T. montanus* Beier) (Atemnidae) - 4100 m (Kilimanjaro)

*Neobisium alticola* Beier (Neobisiidae) - 4100 m (Anatolia)

*Orochernes nepalensis* Beier (Chernetidae) - 4000 m (Nepal)

*Dactylochelifer macrotuberculatus* Krumpal (Cheliferidae) - 4000 m (Nepal)

"*Chelifer*" *baltistanus* di Caporiacco (Cheliferidae) - 3950 m (Karakorum)

*Nudochernes crassus* Beier (Chernetidae) - 3700 m (Elgon)

*Gobichelifer chelanops* (Redikorzev) (Cheliferidae) - 3650 m (Karakorum, Kirgizstan)

*Dactylochelifer popovi* Redikorzev (Cheliferidae) - 3650 m (Tajikistan)

*Calocheiridius crassifemoratus* Beier (Olpiidae) - 3650 m (Elgon)

*Centrochthonius kozlovi* (Redikorzev) (Chthoniidae) - 3650 m (Nepal)

*Megachernes himalayensis* (Ellingsen) (Chernetidae) - 3650 m (Nepal)

*Apocheiridium rossicum* Redikorzev (Cheiridiidae) - 3600 m (Nepal)




Fig. 7. Genera of Pseudoscorpionida living at or above 4000 m in the Old World. A - Hyidae (1); B - Neobisiidae (2); C - Atemnidae (1); D - Cheliferidae (3); E - Chernetidae (3)

*Lechytiya himalayensis* Beier (Lechytiidae) - 3600 m (Nepal)  
*Megachernes soricicola* Beier (Chernetidae) - 3550 m (Nepal)  
*Withius somalicus* Beier (Withiidae) - 3500 m (Elgon)  
*Stenohya gruberi* (Curčić) (Hyidae) - 3500 m (Nepal)  
*Tyrannochthonius rahmi* Beier (Chthoniidae) - 3500 m (Nepal)  
*Tyrannochthonius robustus* Beier (Chthoniidae) - 3500 m (Sichuan, China)  
*Hysterochelifer nepalensis* Beier (Cheliferidae) - 3500 m (Nepal)  
*Nudochernes montanus* Beier (Chernetidae) - 3500 m (Elgon)  
*Nudochernes robustus* Beier (Chernetidae) - 3500 m (Elgon)  
*Rhacochelifer cf. subsimilis* Vachon (Cheliferidae) - 3500 m (Tibesti)

As we can see from this list, from the 25 species 12 are known from Nepal/Tibet, 6 from Elgon/Kilimanjaro, 3 from Karakorum, 1 from Sichuan, 1 from Tajikistan, 1 from Tibesti and 1 from Anatolia. Ten families are represented. Beyond 4000 m we can find only 7 species, and only 1 or 2 to go higher than 4500 m, reaching 5000 m in Nepal.

## References

- BEIER M. 1928. Die Pseudoscorpione des Wiener Naturhistorischen Museums. I. - Ann. Naturhist. Mus. Wien, 42: 285-314.
- BEIER M. 1929. Die Pseudoscorpione des Wiener Naturhistorischen Museums. II. - Ann. Naturhist. Mus. Wien, 43: 341-367.
- BEIER M. 1931. Neue Pseudoscorpione der U.O. Neobisinea. - Mitt. zool. Mus. Berlin, 17: 299-318.
- BEIER M. 1935. Arachnida. I. Pseudoscorpionidea. - In: Mission scientifique de l'Omo, II(6). Mém. Mus. Hist. Nat. Paris, 2: 117-129.
- BEIER M. 1939. The Pseudoscorpionida collected by Percy Sladen Trust Exped. to lake Titikaka. - Ann. Mag. Nat. Hist., 11 (3): 288-290.
- BEIER M. 1943. Neue Pseudoscorpione aus West-, Zentral- und Ostasien. - Ann. Naturhist. Mus. Wien, 53 (2): 74-81.
- BEIER M. 1944. Über Pseudoscorpioniden aus Ostafrika. - Eos, Madrid, 20: 173-212.
- BEIER M. 1951a. On some Pseudoscorpionidea from Kilimanjaro. - Ann. Mag. Nat. Hist., 12 (4): 606-609.
- BEIER M. 1951b. Die Pseudoscorpione Indochinas. - Mém. Mus. Nat. Hist. Nat., Paris (n.s.) 1: 47-123.
- BEIER M. 1953. Eine neue *Neobisium* - Art (Pseudoscorpionidea) aus der Dauphiné. - Ann. Naturhist. Mus. Wien, 59: 155-156.
- BEIER M. 1954. Ein neuer Ophiide (Pseudoscorpionidea) aus dem Hochlande von Peru. - Senckenbergiana, 34 (4-6): 325-326.
- BEIER M. 1955. Pseudoscorpionidea, gesammelt während der schwedischen Expeditionen nach Ostafrika 1937-38 und 1948. - Arkiv för Zoologi, ser. 2, 25 (7): 527-558.
- BEIER M. 1957. Pseudoscorpione gesammelt von Dr K. Lindberg 1956. - Förhandl. Kungl. Fys. Sällsk. I Lund, 27: 145-151.
- BEIER M. 1958. The Pseudoscorpionidea (False-scorpions) of Natal and Zululand. - Ann. Natal Mus., 14 (2): 155-187.
- BEIER M. 1959a. Ergänzungen zur iberischen Pseudoscorpioniden - Fauna. - Eos, Madrid, 35 (2): 113-131.
- BEIER M. 1959b. Zur Kenntnis der Pseudoscorpionidenfauna Afghanistans. - Zool. Jahrb. Syst., Jena, 87 (3): 257-282.



- BEIER M. 1959d. Zur Kenntnis der Pseudoscorpioniden - Fauna des Andengebietes. - Beitr. z. Neotropischen Fauna, 1 (3): 185-228.
- BEIER M. 1959c. Ein neuer *Allochernes* (Pseudoscorpionidea) aus dem Karakorum Gebirge. - Ann. Naturhist. Mus. Wien, 63: 407-408.
- BEIER M. 1959d. Pseudoscorpione aus dem Belgischen Congo gesammelt von Herrn N. Leleup. - Ann. Mus. Congo Belge, Sciences zoologiques, 72: 5-69.
- BEIER M. 1960. Pseudoscorpionidea. Contribution à l'étude de la faune d'Afghanistan 56. - Kungl. Fys. Sällsk. i Lund Forh., 31 (1): 1-4.
- BEIER M. 1961. Nochmals über iberische und marokkanische Pseudoscorpione. - Eos, Madrid, 37 (1): 21-39.
- BEIER M. 1962. Pseudoscorpionidea. - In: Mission Zoologique de l'I.R.S.A.C. en Afrique orientale VIII. - Ann. Mus. Roy. Afrique Centr., Tervuren, sér. in 8, Sci. Zool., 107: 9-37.
- BEIER M. 1964. Pseudoscorpione aus dem Bucegi - Gebirge in Rumanien. - Zool. Anz., 173 (3): 210-212.
- BEIER M. 1966. Die Pseudoscorpioniden der Salomon - Inseln. - Ann. Naturhist. Mus. Wien, 69: 133-159.
- BEIER M. 1967. Pseudoscorpione aus dem tropischen Ostafrika (Kenya, Tansania, Uganda, etc.). - Ann. Naturhist. Mus. Wien, 70: 73-93.
- BEIER M. 1968. Ein neues Chernetiden - Genus (Pseudoscorpionidea) aus Nepal. - Khumbu Himal, 3(1): 17-18.
- BEIER M. 1969. Weitere Beiträge zur Kenntnis der Pseudoscorpione Anatoliens. - Naturhist. Mus. Wien, 73: 189-193.
- BEIER M. 1973a. Beiträge zur Pseudoscorpioniden - Fauna Anatoliens. - Fragmenta entom., Roma, 8 (5): 223-236.
- BEIER M. 1973b. Pseudoscorpionidea von Ceylon. - Ent. Scand. Suppl., 4: 39-55.
- BEIER M. 1974a. Pseudoscorpione aus Sündindien des Naturhistorischen Museums in Genf. - Rev. suisse Zool., 81 (4): 999-1017.
- BEIER M. 1974b. Pseudoscorpione aus Nepal. - Senckenbergiana biol., 55 (4-6): 261-280.
- BEIER M. 1976a. Ergebnisse der Bhutan - Expedition 1972 des Naturhistorischen Museums Basel. Pseudoscorpionidea. - Verhandl. Naturf. Ges. Basel, 85 (1-2): 95 -100.
- BEIER M. 1978. Pseudoscorpione aus Kashmir und Ladakh (Arachnida). - Senckenbergiana biol., 58 (5-6): 415-417.
- BEIER M. 1982. Zoological Results of the British Speleological Expedition to Papua New Guinea 1975. 9. Pseudoscorpionidea. - Acta zool. bulgarica, 19: 43-45.
- BEIER M., H. FRANZ. 1954. 16. Ordnung Pseudoscorpionidea. - In: Franz, H.: Die Nordostalpen im Spiegel ihrer Landtierwelt, 1: 453-459.
- BERON P. 1999. Composition and Biodiversity of the High Mountain Terrestrial Fauna in Bulgaria. - Historia naturalis bulgarica, 10: 13-33.
- BERON P. 2000. Non-insect Arthropoda (Isopoda, Arachnida and Myriapoda) on the high mountains of tropical Africa. - In: Rheinwald, G., ed.: Isolated Vertebrate Communities in the Tropics. Proc. 4<sup>th</sup> Int. Symp., Bonn, Bonn. Zool. Monogr., 46: 153-188.
- CAPORIACCO L. di. 1927. Secondo saggio sulla fauna aracnologica della Carnia e regioni limitrofe. - Mem. Soc. Entom. It., 5: 70-130.
- CHAMBERLIN J. C. 1930. A Synoptic Classification of the False Scorpions or Chelaspinners, with a Report on a Cosmopolitan Collection of the same. - Part II. The Diplosphyronida (Arachnida - Chelonethida). - Ann. Mag. Nat. Hist., ser. 10, 5: 585-620.
- CURČIĆ B. 1980. Pseudoscorpions from Nepal. - Bull. Mus. Hist. Nat. Belgrade, B, 35: 77-101.
- CURČIĆ B. 1984. The genus *Neobisium* Chamberlin, 1930 (Neobisiidae, Pseudoscorpiones, Arachnida): on new species from the USSR and the taxonomy of its subgenera. - Bull. Mus. Hist. Nat., Belgrade, B, 39: 123-153.
- CURČIĆ B. 1985. A revision of some species of *Microcreagra* Balzan, 1892 (Neobisiidae, Pseudoscorpiones) from the USSR and adjacent regions. - Bull. Br. Arachnol. Soc., 6 (8): 331-352.



- DASHDAMIROV S. 1990. A fauna and zoogeography of pseudoscorpiones of the Azerbaijan (Arachnida, Pseudoscorpiones). - In: Fauna and Ecology of Spiders, Scorpions and Pseudoscorpions of the USSR, Proc. Zool. Inst. Leningrad, 226: 105-110.
- DASHDAMIROV S., W. SCHAWALLER. 1992a. Pseudoscorpions of the Caucasian fauna (Arachnida Pseudoscorpionida). - Arthropoda Selecta, 1 (4): 31-72.
- DASHDAMIROV S., W. SCHAWALLER. 1992b. Pseudoscorpions from Middle Asia, Part 1 (Arachnida: Pseudoscorpiones). - Stuttgarter Beitr. Naturk., A, 474:1-18.
- DASHDAMIROV S., W. SCHAWALLER. 1993a. Pseudoscorpions from Middle Asia, Part 2 (Arachnida: Pseudoscorpiones). - Stuttgarter Beitr. Naturk., A, 496:1-14.
- DASHDAMIROV S., W. SCHAWALLER. 1993b. Pseudoscorpions from Middle Asia, Part 3 (Arachnida: Pseudoscorpiones). - Stuttgarter Beitr. Naturk., A, 497: 1-16.
- DASHDAMIROV S., W. SCHAWALLER. 1995. Pseudoscorpions from Middle Asia, Part 4 (Arachnida: Pseudoscorpiones). - Stuttgarter Beitr. Naturk., A, 522: 1-24.
- HARVEY M.S. 1991. Catalogue of the Pseudoscorpionida. - Manchester Univ. Press, Manchester: 726 p.
- HARVEY M. S. 1992. The Phylogeny and Classification of the Pseudoscorpionida (Chelicerata: Arachnida). - Invert. Taxon., 6: 1373-1435.
- HEURTAULT J. 1970. Pseudoscorpions du Tibesti (Tchad). I. Olpiidae. - Bull. Mus. Nat. Hist. Nat. Paris, 41 (2): 1164 -1174.
- HEURTAULT J. 1971. Pseudoscorpions de la région du Tibesti (Sahara méridionale). IV. Cheliferidae. - Bull. Mus. Nat. Hist. Nat., Paris (2), 42: 685-707.
- JANETSCHKEK H. 1949. Tierische Successionen auf hochalpinen Neuland. - Schlern - Schriften, Innsbruck, 67: 1-215.
- JANETSCHKEK H. 1957a. Zoologische Ergebnisse einer Studienreise in die spanische Sierra Nevada (Vorläufige Mitteilung). - Publ. Inst. Biol. Aplic., Barcelona, 26: 135-153.
- JANETSCHKEK H. 1957b. Zur Landtierwelt der Dolomiten. - Der Schlern, 31: 71-86.
- KISHIDA K. 1966. On the altitudinal distribution of the Chelonethida in Japan. - Acta Arachnologica, 20: 6-8.
- KOBACHIDZE D. 1960. Materialien zur Höhenstufenverbreitung der Pseudoscorpionidea in der georgischen SSR. - Zeitschr. der Arbeitsgemeinschaft österr. Entomologen, 12 (2): 103-106.
- KOFLER A. 1972. Die Pseudoscorpione Osttirols. - Mitt. Zool. Ges. Braunau, 1: 286-289.
- KRUMPÁL M. 1987. Ein neuer *Dactylochelifer* aus Nepal Himalaya (Arachnida, Pseudoscorpiones). - Acta Entom. Bohemoslov., 84 (3): 221-226.
- LAZZERONI G. 1969. Contributo alla conoscenza degli Pseudoscorpioni della Regione Veronese (Ricerche sugli Pseudoscorpioni. IV.) - Mem. Mus. Civ. St. Nat. Verona, 16 (1968): 397-418.
- LESSERT R. de. 1911. Pseudoscorpions. - Catalogue des Invertébrés de la Suisse, 5: 1-50.
- MAHNERT V. 1977. Pseudoskorpione (Arachnida) aus dem Tien-Shan. - Ber. nat. - med. Ver. Innsbruck, 64: 89-95.
- MAHNERT V. 1981. Die Pseudoskorpione (Arachnida) Kenyas. I. Neobisiidae und Ideoroncidae. - Revue suisse Zool., 88 (2): 535-559.
- MAHNERT V. 1982a. Die Pseudoskorpione (Arachnida) Kenyas II. Feaellidae; Cheiridiidae. - Revue suisse Zool., 89 (1): 115-134.
- MAHNERT V. 1982b. Die Pseudoskorpione (Arachnida) Kenyas, IV. Garypidae. - Ann. Hist.-Nat. Mus. Nat. Hungarici, 74: 307-329.
- MAHNERT V. 1982c. Die Pseudoskorpione (Arachnida) Kenyas V. Chernetidae. - Revue suisse Zool., 89 (3): 691-712.
- MAHNERT V. 1983a. Die Pseudoskorpione Kenyas VI. Dithidae (Arachnida). - Rev. Zool. afr., 97 (1): 141-157.
- MAHNERT V. 1983b. Die Pseudoskorpione (Arachnida) Kenyas VII. Miratemnidae und Atemnidae. - Revue suisse Zool., 90 (2): 357-398.

- MAHNERT V. 1985. Die Pseudoskorpione (Arachnida) Kenyas. VIII. Chthoniidae. - *Revue suisse Zool.*, 92 (4): 823-843.
- MAHNERT V. 1988. Die Pseudoskorpione (Arachnida) Kenyas. Familien Withiidae und Cheliferidae. - *Tropical Zoology*, 1: 39-89.
- MAHNERT V. 1997. New species and records of Pseudoscorpions (Arachnida, Pseudoscorpiones) from the Canary Islands. - *Rev. suisse Zool.*, 104 (3): 559-585.
- MANI M. S. 1959. On a collection of high altitude scorpions and pseudoscorpions (Arachnida) from the Northwest Himalaya. - *Agra Univ. J. Res. (Sci.)*, 8 (1): 11-16.
- MORIKAWA K. 1968. On some pseudoscorpions from Rolwaling Himal. - *J. College Arts Sci. Chiba Univ.*, 2: 259-263.
- PALMGREN P. 1973. Über die Biotopverteilung waldbodenlebender Pseudoscorpionidea (Arachnoidea) in Finnland und Österreich. - *Commentationes Biologicae, Helsinki*, 61: 1-11.
- REDIKORZEV V. 1918. Pseudoscorpions nouveaux. I. - *Ezhagodnik Zool. Muzeya*, 22: 91-101.
- REDIKORZEV V. 1928. Beiträge zur Kenntnis der Pseudoscorpionenfauna Bulgariens. - *Mitt. Königl. Natuwiss. Inst. Sofia*, 1: 118-141.
- REDIKORZEV V. 1949. [Pseudoscorpionidea of Central Asia]. - *Trav. Inst. Zool. Acad. Sci. U.R.S.S.*, 8: 638-668 (in Russian).
- SATO H. 1980. Altitudinal Distribution of Soil Pseudoscorpions on Mt. Chokai. - *Edaphologia*, 22: 9-14.
- SCHAWALLER W. 1983a. Pseudoskorpione aus dem Kaukasus (Arachnida). - *Stuttgarter Beitr. Naturk.*, A, 362: 1-24.
- SCHAWALLER W. 1983b. Neue Pseudoskorpion - Funde aus dem Nepal-Himalaya (Arachnida: Pseudoscorpionidea). - *Senckenbergiana biol.*, 63: 105-111.
- SCHAWALLER W. 1983c. Pseudoskorpione aus dem Norden des Iran (Arachnida: Pseudoscorpionidea). - *Stuttgarter Beitr. Naturk.*, A, 385: 1-12.
- SCHAWALLER W. 1985. Pseudoskorpione aus der Sowjetunion (Arachnida: Pseudoscorpiones). - *Stuttgarter Beitr. Naturk.*, A, 385: 1-12.
- SCHAWALLER W. 1986. Pseudoskorpione aus der Sowjetunion, Teil 2 (Arachnida: Pseudoscorpiones). - *Stuttgarter Beitr. Naturk.*, A, 385: 1-12.
- SCHAWALLER W. 1987. Neue Pseudoskorpion-Funde aus dem Nepal-Himalaya, II (Arachnida: Pseudoscorpiones). - *Senckenbergiana biol.*, 68 (1-3): 199-221.
- SCHAWALLER W. 1988. Neue Pseudoskorpion - Funde aus dem Nepal-Himalaya, II. (Arachnida: Pseudoscorpiones). - *Stuttgarter Beitr. Naturk.*, A, 396: 1-15.
- SCHAWALLER W. 1989. Pseudoskorpione aus der Sowjetunion, Teil 3 (Arachnida: Pseudoscorpiones). - *Stuttgarter Beitr. Naturk.*, A, 440: 1-30.
- SCHAWALLER W. 1991. Neue Pseudoskorpion - Funde aus dem Nepal-Himalaya, III. (Arachnida: Pseudoscorpiones). - *Revue suisse Zool.*, 98 (4): 769-789.
- SCHAWALLER W. 1994. Pseudoskorpione aus Thailand (Arachnida: Pseudoscorpiones). - *Revue suisse Zool.*, 101 (3): 725-759.
- SCHAWALLER W. 1995. Review of the Pseudoscorpion Fauna of China (Arachnida: Pseudoscorpionida). - *Rev. suisse Zool.*, 102 (4): 1045-1064.
- SCHAWALLER W., S. DASHDAMIROV. 1988. Pseudoskorpion - Funde aus dem Kaukasus, Teil 2 (Arachnida). - *Stuttgarter Beitr. Naturk.*, A, 415: 1-51.
- SCHMÖLZER K. 1962. Die Kleintierwelt der Nunatakter als Zeugen einer Eiszeitüberdauerung. - *Mitt. zool. Mus. Berlin*, 38 (2): 171-400.
- THALER K. 1979. *Fragmenta Faunistica Tirolensia*, IV (Arachnida: Acari: Caeculidae; Pseudoscorpiones; Scorpiones; Opiliones; Aranei. Insecta: Dermaptera; Thysanoptera; Diptera Nematocera: Mycetophilidae, Psychodidae, Limoniidae und Tipulidae). - *Ver. Mus. Ferdinandeum*, 59: 49-83.
- TULLGREN A. 1910. Pedipalpi, Scorpiones, Solifugae, Chelonethi. - In: *Sjöstedt, Wiss. Erg. Kilimandjaro-Meru Expedition, Stockholm* 3: 1-15.

- VACHON M. 1945. Mission scientifique de l'Omo, vol.6, part 10. Chernètes. - Mém. Mus. Nat. Hist. Nat., Paris, n.s., 19: 187-197.
- WÜRMLI M. 1972. Zur vergleichenden Synökologie und Faunistik der kryptozoischen Makroarthropoden Mitteleuropas und Südtaliens. - Mitt. Schweiz. Ent. Ges., 45 (1-3): 157-216.

Received on 01.09.2001

Author's address:  
Dr Petar Beron  
National Museum of Natural History  
Tsar Osvoboditel Blvd. 1  
1000 Sofia, Bulgaria

## Върху високoplanинските псевдоскорпиони (Arachnida: Pseudoscorpionida) на Стария свят

Петър БЕРОН

(Резюме)

От известните около 435 рода псевдоскорпиони в света от 24 семейства, 66 рода и общо 158 вида са познати от Стария свят на височина над 2200 m. Те спадат към 16 семейства: Chthoniidae (17 вида), Lechytiidae (2 вида), Tridenchthoniidae (7 вида), Geogarypidae (8 вида), Olpiidae (8 вида), Hyidae (4 вида), Gymnobisiidae (1 вид), Ideoroncidae (1 вид), Neobisiidae (25 вида), Syarinidae (2 вида), Cheiridiidae (5 вида), Sternophoridae (1 вид), Atemnidae (14 вида), Cheliferidae (18 вида), Chernetidae (37 вида), Withiidae (8 вида). В Европа само 4 вида достигат 3000 m: *Neobisium jugorum* L. Koch (Алпите, 3600 m), *N. nivale* Beier (Сиера Невада, 3481 m), *N. anatolicum* Beier (Кавказ, 3000 m) и *N. carcinoides* Hermann (Алпите, 3000 m, Пирин, 2914 m). В планините на Централна и Източна Африка над 2200 m живеят най-малко 26 вида, от които 13 достигат 3000 m, 4 - 3500 m и само *Titanatemnus palmquisti* Tulgren надминава 4000 m (до 4100 m в Танзания и Кения). В Хималаите неопределен до вид псевдоскорпион от сем. Hyidae или Neobisiidae (juv.) е намерен на височина около 5000 m, която е и максималната за представителите на този разред в света.

В Хималаите над 2200 m са намерени 29 вида псевдоскорпиони, над 3000 m - 17, над 4000 m - 3 (*Stenohya martensi* Schawaller - 4700 m, *Orochernes nepalensis* Beier и *Dactylochelifer macrotuberculatus* Krumpal, 4000 m, всички от Непал). Видовете над 2200 m спадат към 20 рода и 11 семейства. Интересен факт е, че високoplanинските псевдоскорпиони в Източна и Централна Африка също спадат към 11 семейства, десет от които са общи за двата района.

От планините на Централна Азия са известни не по-малко от 12 вида псевдоскорпиони от 8 семейства, обитаващи райони над 2200 m, включително 11 вида над 3000 m и 4 над 3500 m. Шампиони са *Bisetocreagris kaznakovi* (Redikorzev) - 4810 m (Neobisiidae, Тибет), *Dactylochelifer brachialis* Beier (4200 m, Каракорум), *Gobichelifer chelanops* (Redikorzev) (3650 m, Каракорум) и "*Chelifer*" *baltistanus* di Saporiasso (3950 m, Каракорум) (и четирите спадат към сем. Cheliferidae), докато всички европейски псевдоскорпиони над 3000 m принадлежат към сем. Neobisiidae. В Европа никой от многобройните видове Cheliferidae не живее над 2200 m.